

## Transient analysis of stochastic fluid models

B. Sericola<sup>1</sup>

*IRISA-INRIA, Campus de Beaulieu, 35042 Rennes Cédex, France*

Received 28 May 1997; received in revised form 19 December 1997

---

### Abstract

We analyze the transient behavior of stochastic fluid flow models in which the input and output rates are controlled by a finite homogeneous Markov process. Such models are used in asynchronous transfer mode (ATM) to evaluate the performance of fast packet switching and in manufacturing systems for the performance of producers and consumers coupled by a buffer. The transient analysis of such models has already been considered in earlier works and solutions have been obtained by the use of Laplace transform. We derive in this paper a new transient solution only based on recurrence relations. We show that this solution is particularly interesting for its numerical properties. The limiting behavior of the solution is also considered. We empirically show that the algorithm for computing the transient solution can be stopped when some stationary behavior is detected. © 1998 Elsevier Science B.V.

*Keywords:* ATM; Transient analysis; Fluid flow models; Markov modulated rate process; Buffer content

---

### 1. Introduction

A stochastic fluid flow model describes the behavior of a fluid level in a storage device. The input and output rates are supposed to be controlled by a finite homogeneous Markov process. Such models are used in asynchronous transfer mode (ATM) to evaluate the performance of fast packet switching and in manufacturing systems for the performance of producers and consumers coupled by a buffer. There is a large number of papers dealing with the analysis of stochastic fluid flow models. Most of these papers consider such models in stationary regime. Anick et al. [1] and Kosten [6] analyzed the fluid model for several on–off input sources controlled by a two-state homogeneous Markov process. Mitra [7,8] generalizes this model by considering multiple on–off inputs and outputs. In [14] Stern and Elwalid considered such models for separable Markov modulated rate process which led to a solution of the equilibrium equations expressed as a sum of terms in Kronecker product form. In [4] Igel'nik et al. derive a new approach, based on the use of interpolating polynomials, for the computation of the buffer overflow probability. An extensive list of references can be found in [12,13].

---

<sup>1</sup> E-mail: bruno.sericola@irisa.fr.

For what concerns the transient analysis stochastic fluid flow models controlled by a finite Markov process. Narayanan and Kulkarni [10] derive explicit expressions for the Laplace transform of the joint distribution for the first time the buffer becomes empty and the state of the Markov process at that time. Guillemin et al. consider the unbuffered model in [3] and obtain a method to compute transient characteristics, such as the congestion period, with an unbounded number of exponential on–off sources. These results have been extended by Dupuis et al. in [2] to the case where the off periods are phase-type.

The Laplace transform has been largely used to evaluate the transient behavior of fluid flow models. In [11] Ren and Kobayashi studied the transient distribution of the buffer content for exponential on–off sources of a single type. The same authors deal with the case of multiple types of inputs in [5]. These studies have been extended to the Markov modulated input rate model by Tanaka et al. in [15].

In this paper, we consider a general stochastic fluid flow model in which the buffer is infinite and the input and output rates are controlled by a finite homogeneous Markov process. For this model we derive a new transient solution for the distribution of the buffer content. This solution does not make use of any transform, it is only based on simple recurrence relations which are particularly interesting for their numerical properties. The algorithm implementing this solution is very accurate since it uses essentially nonnegative numbers bounded by one and it gives results with an error tolerance that can be specified in advance. Furthermore, by considering the limiting behavior of the solution, we empirically show that the algorithm can be stopped when some stationary behavior is detected.

The remainder of the paper is organized as follows. We describe in Section 2 the model and we present our new transient solution with some of its properties. In Section 3 we describe the algorithm implementing the solution. We empirically show in Section 4, through numerical examples, that the computation time of the solution can be considerably reduced by considering the limiting behavior of the solution. Section 5 is devoted to some conclusions.

## 2. A new transient solution

We describe in this section a general fluid model with an infinite buffer for which the input and output rates are controlled by a homogeneous Markov process  $X = \{X_s, s \geq 0\}$  on the finite state space  $S$  with infinitesimal generator  $A$  and initial probability distribution  $\alpha$ . The number of states is denoted by  $|S|$ . The amount of fluid in the buffer at time  $t$  is denoted by  $Q_t$  and we suppose that  $Q_0 = 0$ . The pair  $(X_t, Q_t)$  forms a Markov process having a pair of discrete and continuous states. Let  $\rho_i$  be the input rate and  $c_i$  be the output rate when the Markov process  $X$  is in state  $i$ . We denote by  $d_i$  the effective input rate of state  $i$ , that is  $d_i = \rho_i - c_i$ . Let  $m + 1, m < |S|$ , be the number of distinct values among all the effective rates  $d_i$ . These  $m + 1$  distinct effective rates are denoted by  $r_0, r_1, \dots, r_m$  and ordered as follows

$$r_m > r_{m-1} > \dots > r_u > 0 \geq r_{u-1} > \dots > r_1 > r_0,$$

where  $u$  is the index of the smallest positive effective rate. The state space  $S$  of the process  $X$  can then be divided into  $m + 1$  disjoint subsets  $B_m, B_{m-1}, \dots, B_0$  where  $B_i$  is composed by the states  $i$  of  $S$  having the same effective rate  $r_i$ , that is  $B_i = \{j \in S | d_j = r_i\}$ . We will denote by  $|B_i|$  the cardinal of subset  $B_i$ .

For a fixed  $t > 0$ , the random variable  $Q_t$  takes its values in the interval  $[0, r_m t]$ . For  $t > 0$ , the distribution of  $Q_t$  has  $m - u + 1$  jumps at positive values and one jump at point 0 corresponding to the case where the buffer is empty at time  $t$ . The jumps at the  $m - u + 1$  positive values correspond to the case where the Markov process  $X$  remains during the whole interval  $[0, t]$  in the different subsets  $B_u, B_{u+1}, \dots, B_m$ ,

provided that the initial probabilities of these subsets are positive. These jump, probabilities are then given, for  $j = u, u + 1, \dots, m$ , by

$$\Pr\{X_t = i, Q_t = r_j t\} = \alpha_{B_j} e^{A_{B_j B_j} t} \mathbf{1}_i \quad \text{if } i \in B_j,$$

where  $A_{B_j B_j}$  is the sub-infinitesimal generator of dimension  $|B_j|$  obtained from  $A$  by considering only the internal transitions of the subset  $B_j$  and  $\alpha_{B_j}$  is the subvector of dimension  $|B_j|$  obtained from the row vector  $\alpha$  by considering the initial probabilities of the subset  $B_j$ . The vector  $\mathbf{1}_i$  is the column vector whose  $i$ th entry is 1 and the others 0, its dimension being given by the context ( $|B_j|$  in this relation).

The jump at point 0 is not so easy to obtain since the process  $X$  can eventually visit all the subsets  $B_i$  before the buffer becomes empty at time  $t$ .

Let  $F_i(t, x) = \Pr\{X_t = i, Q_t > x\}$ . We then have the following partial differential equation, see for instance [15]:

$$\frac{\partial F_i(t, x)}{\partial t} = -d_i \frac{\partial F_i(t, x)}{\partial x} + \sum_{r \in S} F_r(t, x) A(r, i). \tag{1}$$

We denote by  $P$  the transition probability matrix of the uniformized Markov chain associated to  $X$  and by  $\lambda$  the uniformization rate which verifies  $\lambda \geq \max(-A(i, i), i \in S)$ . The matrix  $P$  is then related to  $A$  by  $P = I + A/\lambda$ , where  $I$  denotes the identity matrix. In the following, to simplify notation, we will consider  $X$  as the uniformized process. For every  $i, j = 0, \dots, m$ , we denote by  $P_{B_i B_j}$  the submatrix of  $P$  containing the transition probabilities from states of  $B_i$  to states of  $B_j$  and for  $n \geq 1$  we denote by  $(\alpha P^n)_{B(i)}$  the subvector of the row vector  $\alpha P^n$  containing the entries corresponding to the subset  $B_i$ .

The main result of this paper, which is the distribution of the pair  $(X_t, Q_t)$  is given by the following theorem.

**Theorem 1.** *For every  $i \in S$ , we have*

$$F_i(t, x) = \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b_i^{(j)}(n, k), \tag{2}$$

where  $x_j = (x - r_{j-1}^+ t) / (r_j - r_{j-1}^+ t)$  if  $x \in [r_{j-1}^+ t, r_j t)$ , for  $j = u, u + 1, \dots, m$ , with  $r_{j-1}^+ = 0$  for  $j = u$  and  $r_{j-1}^+ = r_{j-1}$  for  $j > u$ . The coefficients  $b_i^{(j)}(n, k)$  are given by the following recursive expressions on the row vectors  $b_{B_l}^{(j)}(n, k) = (b_i^{(j)}(n, k))_{i \in B_l}$  for  $0 \leq l \leq m$  and  $u \leq j \leq m$ .

- for  $j \leq l \leq m$  :
  - for  $n \geq 0$ :  $b_{B_l}^{(j)}(n, 0) = (\alpha P^n)_{B_l}$  and  $b_{B_l}^{(j)}(n, 0) = b_{B_l}^{(j-1)}(n, n)$  for  $j > u$ ,
  - for  $1 \leq k \leq n$ :

$$b_{B_l}^{(j)}(n, k) = \frac{r_l - r_j}{r_l - r_{j-1}^+} b_{B_l}^{(j)}(n, k - 1) + \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k - 1) P_{B_i B_l},$$

- for  $0 \leq l \leq j - 1$ :
  - for  $n \geq 0$ :  $b_{B_l}^{(j)}(n, n) = 0_{B_l}$  and  $b_{B_l}^{(j)}(n, n) = b_{B_l}^{(j+1)}(n, 0)$  for  $j < m$ ,

- for  $0 \leq k \leq n - 1$ :

$$b_{B_l}^{(j)}(n, k) = \frac{r_{j-1}^+ - r_l}{r_j - r_l} b_{B_l}^{(j)}(n, k + 1) + \frac{r_j - r_{j-1}^+}{r_j - r_l} \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k) P_{B_i B_l}.$$

**Proof.** See Appendix A.  $\square$

Formula (2) is particularly interesting from a computational point of view. Indeed, for every  $j = u, \dots, m$  and  $x \in [r_{j-1}^+ t, r_j t]$  we have  $x_j \in [0, 1]$ ,

$$0 \leq \frac{r_l - r_j}{r_l - r_{j-1}^+} = 1 - \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \leq 1 \quad \text{for } l = j, \dots, m,$$

and

$$0 \leq \frac{r_{j-1}^+ - r_l}{r_j - r_l} = 1 - \frac{r_j - r_{j-1}^+}{r_j - r_l} \leq 1 \quad \text{for } l = 0, \dots, j - 1.$$

It is then easy to check that for every  $i \in S, j = u, \dots, m, n \geq 0$  and  $k = 0, \dots, n$  we have  $b_i^{(j)}(n, k) \in [0, 1]$ . Moreover the error truncation of the series in (2) can be determined in advance. These properties are very important for what concerns the numerical stability of the computation.

For a given error tolerance  $\varepsilon$ , we define integer  $N$  as

$$N = \min \left\{ n \in \mathbb{N} \mid \sum_{i=0}^n e^{-\lambda t} \frac{(\lambda t)^i}{i!} \geq 1 - \varepsilon \right\}.$$

We then get, for every  $i \in S$ ,

$$F_i(t, x) = \sum_{n=0}^N e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b_i^{(j)}(n, k) + e(N),$$

where the rest of the series  $e(N)$  satisfies  $e(N) \leq \varepsilon$ .

The main computational effort is due to the computation of the  $b_{B_l}^{(j)}(n, k)$  given in Theorem 1. To illustrate the recurrence relation, we proceed as done in [9] for the performability computation. For each  $j = u, \dots, m$ , we define a partition of the state space  $S$  as

$$U_j = B_m \cup \dots \cup B_j \quad \text{and} \quad D_j = B_{j-1} \cup \dots \cup B_0,$$

and denoting by  $T$  the transpose operator, we also define the following column vectors:

$$b_{U_j}(n, k) = (b_{B_m}^{(j)}(n, k), \dots, b_{B_j}^{(j)}(n, k))^T$$

$$b_{D_j}(n, k) = (b_{B_{j-1}}^{(j)}(n, k), \dots, b_{B_0}^{(j)}(n, k))^T.$$

With this notation, Fig. 1 illustrates the sequence of computations (drawn only for  $n = 0, 1, 2, 3$ ) that have to be done in order to evaluate the  $b_{B_l}^{(j)}(n, k)$ 's. The upper (resp. lower) part of cell  $(n, k)$  in triangle  $j$




Fig. 1. In cell  $(n, k)$  the vectors  $b_{U_j}(n, k)$  and  $b_{D_j}(n, k)$ .




Fig. 2. Computation of cell  $(n, k)$ .

contains the vector  $b_{U_j}(n, k)$  (resp.  $b_{D_j}(n, k)$ ). The computation is done in a line by line manner over all the triangles following the arrows in Fig. 1. Note that the upper part of the diagonal of each triangle of cells is reported in the upper part of the first column of the next one and the lower part of the first column of each triangle of cells is reported in the lower part of the diagonal of the previous triangle of cells. The starting points are given, for  $j = u, \dots, m$  and for every  $n \geq 0$ , by

$$b_{U_j}(0, 0) = (\alpha_{B_m}, \dots, \alpha_{B_j})^T, \quad b_{D_j}(0, 0) = (0_{B_{j-1}}, \dots, 0_{B_0})^T,$$

and

$$b_{U_u}(n, 0) = ((\alpha P^n)_{B_m}, \dots, (\alpha P^n)_{B_j})^T, \quad b_{D_m}(n, n) = (0_{B_{m-1}}, \dots, 0_{B_0})^T.$$

The way in which the computation of each cell  $(n, k)$  is performed is shown in Fig. 2. It is now easy to evaluate the complexity of this method. The computation of one cell consists essentially in a vector matrix product. If  $d$  denotes the maximum number of nonzero entries in each column of the matrix  $P$ , the computational complexity of a cell is  $O(d|S|)$ . There are  $m-u+1$  triangles each containing  $(N+1)(N+2)/2$  cells. The computational complexity of our method is then  $O(d|S|(m-u+1)N^2/2)$ . We see from Figs. 2

and 1 that it is sufficient to store two rows of cells in order to compute the  $b_i^{(j)}(n, k)$ . Thus the storage complexity of our method is  $O((m - u + 1)N|S|)$ .

### 3. Stationarity detection

We empirically show in this section that the algorithm described above can be stopped when the stationary behavior of the model is detected.

Let us denote by  $\pi$  the stationary distribution of the Markov process  $X$ . We suppose that the stability condition is satisfied, that is

$$\rho = \frac{\sum_{i \in S} \rho_i \pi_i}{\sum_{i \in S} c_i \pi_i} < 1,$$

where  $\rho$  is the traffic intensity, so that the limiting behavior exists.

With this assumption, we have for every  $j = u, \dots, m$ ,

$$\lim_{t \rightarrow \infty} \Pr\{Q_t > r_j t\} = 0.$$

From relation (2), we have for every  $j = u, \dots, m$ , and  $x \in [r_{j-1}^+ t, r_j t)$

$$\Pr\{Q_t > x\} = \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b^{(j)}(n, k),$$

where  $x_j$  is as in Theorem 1 and

$$b^{(j)}(n, k) = \sum_{i \in S} b_i^{(j)}(n, k).$$

The following theorem gives an upper bound of the  $b_i^{(j)}(n, k)$ . If  $v$  and  $w$  are two vectors having the same dimension, the notation  $v \leq w$  means that the inequality stands for each of their entry, that is  $v_i \leq w_i$  for every  $i$ .

**Theorem 2.** For every  $n \geq 0$ , for every  $j = u, \dots, m$ , for every  $0 \leq k \leq n$  and for every  $0 \leq l \leq m$ , we have

$$b_{B_l}^{(j)}(n, k) \leq (\alpha P^n)_{B_l}. \tag{3}$$

**Proof.** See Appendix B.  $\square$

Using this theorem, we easily verify that  $b^{(j)}(n, k) \leq 1$ .

**Theorem 3.** For every  $n \geq 1$ , for every  $j = u, \dots, m$  and for every  $1 \leq k \leq n$ , we have

$$b_{B_l}^{(j)}(n, 0) \leq b_{B_l}^{(j-1)}(n, n) \quad \text{for } j > u, \tag{4}$$

$$b_{B_l}^{(j)}(n, k) \leq b_{B_l}^{(j)}(n, k - 1) \tag{5}$$

**Proof.** See Appendix C.  $\square$

This theorem shows that for fixed  $n$  and  $i \in S$ , the sequences  $b_i^{(j)}(n, k)$  and  $b^{(j)}(n, k)$  are wide-sense decreasing in both  $j$  and  $k$ . It follows that  $b^{(j)}(n, k)$  can be interpreted as the complementary distribution function of a discrete random variable which is the discrete version of  $Q_t$ . For this discrete random variable, the integer  $n$  represents the number of transitions over the interval  $(0, t)$  for the uniformized Markov chain of  $X$ . Thus we conjecture that the limits  $\lim_{n \rightarrow \infty} b^{(j)}(n, k)$  exist and in this case we must have necessarily

$$\lim_{n \rightarrow \infty} b^{(u)}(n, n) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} b^{(j)}(n, 0) = 0 \quad \text{for } j \geq u + 1.$$

The stationarity detection consists in stopping the computation of the numbers  $b_i^{(j)}(n, k)$  when the values  $d^{(u)}(n, 0) = |b^{(u)}(n, 0) - b^{(u)}(n - 1, 0)|$ ,  $b^{(u)}(n, n)$  and  $b^{(j)}(n, 0)$ ,  $j \geq u + 1$ , are sufficiently small. This can be done as follows.

Consider the integer  $N$  defined in Section 2. We define the integer  $N_u$  as

$$N_u = \min\{n \mid 1 \leq n < N \text{ and } d^{(u)}(n, 0) \leq \varepsilon/3 \text{ and } b^{(u)}(n, n) \leq \varepsilon/3\}$$

and for  $j = u + 1, \dots, m$ , we define the integers  $N_j$  as

$$N_j = \min\{n \mid 1 \leq n < N \text{ and } b^{(j)}(n, 0) \leq \varepsilon\}.$$

When  $N_j$  does not exist, we set  $N_j = N$ . If all the  $N_j$  are equal to  $N$ , we obtain the exact solution described in Section 2.

The approximation made here consists in considering that for  $n \geq N_u$ , we have  $|b^{(u)}(n, k) - \lim_{n \rightarrow \infty} b^{(u)}(n, k)| \leq \varepsilon/3$  for  $k = 0, \dots, N_u$  and  $b^{(u)}(n, N_u) \leq \varepsilon/3$  and that for every  $j = u + 1, \dots, m$  and for  $n \geq N_j$ , we have  $b^{(j)}(n, 0) \leq \varepsilon$ .

In practice, we often observe that for  $n \geq N_j$ , the sequence  $b^{(j)}(n, k)$  are wide-sense monotone, so the approximation is justified.

From Theorem 3, we can easily check that we have

$$N_m \leq N_{m-1} \leq \dots \leq N_u,$$

so, when, for  $j > u + 1$ , the integer  $N_j$  is reached, we stop the computation over triangle  $j$  (see Fig. 1) and we set  $b_{D_{j-1}}(N_j + 1, N_j + 1) = 0$ . The computation then continues over triangles  $u, u + 1, \dots, j - 1$ .

We then have for  $j > u + 1$  and  $x \in [r_{j-1}t, r_j t)$ ,

$$\Pr\{Q_t > x\} = \sum_{n=0}^{N_j} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b^{(j)}(n, k) + e(N_j),$$

where the rest of series  $e(N_j)$  satisfies under the approximation hypothesis

$$\begin{aligned} e(N_j) &= \sum_{n=N_j+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b^{(j)}(n, k) \\ &\leq \sum_{n=N_j+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b^{(j)}(n, 0) \\ &\leq \varepsilon \sum_{n=N_j+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \\ &\leq \varepsilon. \end{aligned}$$

For  $j = u$  and  $x \in [0, r_u t)$ , we get

$$\begin{aligned} \Pr\{Q_t > x\} &= \sum_{n=0}^{N_u} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_u^k (1 - x_u)^{n-k} b^{(u)}(n, k) \\ &+ \sum_{n=N_u+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^{N_u} \binom{n}{k} x_u^k (1 - x_u)^{n-k} b^{(u)}(N_u, k) + e(N_u). \end{aligned}$$

The second sum which is infinite can be easily expressed as a finite one; we then obtain

$$\begin{aligned} \Pr\{Q_t > x\} &= \sum_{n=0}^{N_u} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_u^k (1 - x_u)^{n-k} b^{(u)}(n, k) \\ &+ \sum_{k=0}^{N_u} e^{-\lambda t x_u} \frac{(\lambda t x_u)^k}{k!} b^{(u)}(N_u, k) \left[ 1 - \sum_{n=0}^{N_u-k} e^{-\lambda t(1-x_u)} \frac{(\lambda t(1-x_u))^n}{n!} \right] + e(N_u), \end{aligned}$$

where the rest of series  $e(N_u)$  verifies

$$\begin{aligned} e(N_u) &= \sum_{n=N_u+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^{N_u} \binom{n}{k} x_u^k (1 - x_u)^{n-k} (b^{(u)}(n, k) - b^{(u)}(N_u, k)) \\ &+ \sum_{n=N_u+1}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=N_u+1}^n \binom{n}{k} x_u^k (1 - x_u)^{n-k} b^{(u)}(n, k). \end{aligned}$$

Under our approximation hypothesis, we get for  $n \geq N_u$ :

$$|b^{(u)}(n, k) - b^{(u)}(N_u, k)| \leq 2\varepsilon/3 \quad \text{for } k \leq N_u$$

and  $b^{(u)}(n, k) \leq b^{(u)}(n, N_u) \leq \varepsilon/3$  for  $k \geq N_u + 1$ . It follows that we finally obtain  $|e(N_u)| \leq \varepsilon$ .

The complexity of this approximation is now a function of the truncation integers  $N_i$ . The number of cells that must be computed in triangles  $i$  is equal to  $(N_i + 1)(N_i + 2)/2$ . So as for the exact algorithm, we easily obtain the computational complexity of the approximation which is  $O(d|S| \sum_{i=u}^m N_i^2/2)$ . By comparing the computational complexities of the exact and of the approximation method, we see that, the approximation, if sufficiently accurate, must be used for large values of  $m$ . We will see in the next section that the values of the  $N_i$  can be very small with respect to  $N$  with a very high accuracy for the results obtained by the approximation.

#### 4. Numerical examples

We present here some numerical results to illustrate our new solution technique and the approximation based on the stationarity detection.

We consider  $m$  statistically independent and identical on–off sources. For each source, we assume that the on periods and the off periods form an alternating renewal process and their durations are exponentially distributed with mean  $\beta^{-1}$  and  $\gamma^{-1}$ , respectively. When a source is in the state on, it generates packets (or cells in the ATM terminology) at rate  $\theta$ . We denote by  $C$  the multiplexer’s output link capacity. Let  $X_s$  be the number of sources in the state on at time  $s$ . The process  $X = \{X_s, s \geq 0\}$  is then a homogeneous



Markov process over the state space  $S = \{0, 1, \dots, m\}$ . Its infinitesimal generator  $A$  is a tridiagonal matrix whose entries are  $A(i, i - 1) = i\beta$  for  $i = 1, \dots, m$ ,  $A(i, i + 1) = (m - i)\gamma$  for  $i = 0, \dots, m - 1$ , and so  $A(i, i) = -i\beta - (m - i)\gamma$  for  $i = 0, \dots, m$ . For each  $i \in S$ , we have  $\rho_i = i\theta$  and  $c_i = C$ . The traffic intensity  $\rho$  is then

$$\rho = \frac{m\theta\gamma}{C(\beta + \gamma)}.$$

We fix  $\theta = 1$ ,  $\beta = 1$  and  $C = 0.8$ . This gives  $u = 1$  and so the number of triangles that we have to consider is equal to  $m$ . We consider various values of the number  $m$  of sources and of the off rate  $\gamma$  or of the traffic intensity  $\rho$ . The error tolerance is fixed to  $\varepsilon = 10^{-5}$ . Figs. 3–6 have been obtained using the exact algorithm and Fig. 7 has been obtained using the approximation method detecting the stationary behavior of the model.

Fig. 3 shows the complementary distribution of the buffer content at time  $t$  for various values of  $t$ . There are two input sources, the traffic intensity is  $\rho = 5/6$  and both sources are initially in the off state. It can be noted that both distributions for  $t = 100$  and  $t = 200$  are very near to each other, which means that the stationary regime seems to be reached.

Fig. 4 shows the complementary emptiness function  $\Pr(Q_t > 0)$  for 2, 5 and 10 sources when the traffic intensity is  $\rho = 5/6$  and all the input sources are initially in the off state. It can also be noted the convergence of the curves to the traffic intensity  $\rho$ .

Fig. 5 is particularly interesting from a numerical point of view. The value of the time is fixed to  $t = 1$ , the number of input sources is  $m = 2$  and the traffic intensity is  $\rho = 5/6$ . This figure shows the complementary distribution of  $Q_1$  for different initial probability distributions, which correspond to the case where all the input sources are off, that is  $X_0 = 0$ , the case where the input sources are in stationary regime, that is the distribution of  $X_0$  is  $\pi$ , and the case where all the input sources are on, that is  $X_0 = 2$ . When  $X_0 = 0$ , the distribution has only one jump at point 0 and it is not differentiable at point  $x = r_1t = 0.2$ . When the distribution of  $X_0$  is  $\pi$ , we observe the three discontinuities at points  $x = 0$ ,  $x = r_1t = 0.2$  and




Fig. 3. From bottom to the top,  $\Pr\{Q_t > x\}$  versus  $x$  for  $t = 10, 20, 30, 50, 100, 200$ ,  $X_0 = 0$ ,  $m = 2$  and  $\rho = 5/6$ .




Fig. 4. From bottom to the top,  $\Pr\{Q_t > 0\}$  versus  $t$  for  $m = 2, 5, 10$ ,  $X_0 = m$  and  $\rho = 5/6$ .

$x = r_2 t = 1.2$ . These two last discontinuities are easy to determine. For instance, we have  $\Pr\{Q_1 = 0.2\} = \pi_1 e^{-(\beta+\gamma)t} = 4 e^{-1.5}/9$ . The computation of  $\Pr\{Q_1 > 0.2 - 10^{-16}\} - \Pr\{Q_1 > 0.2\}$  using our algorithm gives exactly this result, the precision obtained is greater than  $10^{-10}$ . The same observation holds at point  $x = 1.2$ . Finally when  $X_0 = 2$ , we observe the two jumps at points  $x = 0$  and  $x = r_2 t = 1.2$ . As before, it is easy to check that, at point  $x = r_2 t = 1.2$ , the result obtained using our algorithm is highly accurate. We also observe that the distribution is not differentiable at point  $x = r_1 t = 0.2$ .

Fig. 6 shows the complementary distribution of  $Q_{100}$  for various values of the traffic intensity  $\rho$ , including values greater than 1. The number of input sources is  $m = 2$  and both sources are initially off. For instance, we have  $\Pr\{Q_{100} > 45\} = 0.0001$  for  $\rho = 1.25$ .

Fig. 7 shows the complementary distribution of  $Q_t$  for various values of  $t$ . The traffic intensity is  $\rho = 5/6$ , the number of input sources is  $m = 50$  and all the sources are initially off. This figure has been obtained by using the approximation method based on the stationarity detection. For  $t = 10$ , we obtained for the different truncation steps  $N = 598$ ,  $N_i = 51 - i$  for  $i = 5, \dots, 50$ ,  $N_4 = 374$  and  $N_3 = N_2 = N_1 = N$ . This shows the important gain in computational complexity obtained by the approximation method. To evaluate the accuracy of the approximation method, we have executed the exact algorithm with the same input parameters. We have observed that the greatest difference between the results of the two algorithms is equal to  $2.2 \times 10^{-6}$ . This shows that our approximation method is highly accurate even for small values of  $t$ . For  $t > 10$  we obtain, as expected, an accuracy still higher than for  $t = 10$ .

## 5. Conclusion

We developed a new transient solution of a fluid model with an input and output controlled by a homogeneous Markov process. Our solution does not make use of any transform, as done in previous works.

It is based on simple recurrence relations which are particularly interesting for their numerical properties. The algorithm implementing this solution is very accurate since it uses essentially nonnegative numbers




Fig. 5. From bottom to the top,  $\Pr\{Q_1 > x\}$  versus  $x$  for  $X_0 = 0$ ,  $X_0 \sim \pi$  and  $X_0 = 2$  when  $m = 2$  and  $\rho = 5/6$ .




Fig. 6. From bottom to the top,  $\Pr\{Q_{100} > x\}$  versus  $x$  for  $\rho = 0.75, 1, 1.25, 1.5$ ,  $m = 2$  and  $X_0 = 0$ .

bounded by one and it gives results with an error tolerance that can be specified in advance. We also developed an approximation method based on the detection of the stationary regime of the model. It has been shown through numerical examples that, as the exact method, this approximation method is highly accurate. Moreover its computational time can be very low with respect to the exact method.




Fig. 7. From bottom to the top,  $\Pr\{Q_t > x\}$  versus  $x$  for  $t = 10, 20, 50, 80, 100, 150, 200$  when  $X_0 = 0, m = 50$  and  $\rho = 5/6$ .

**Appendix A. Proof of Theorem 1**

For  $t > 0$  and  $x \in (r_{j-1}^+t, r_j t)$ , for  $j = u, u + 1, \dots, m$ , we write the solution of Eq. (1) for every  $i \in S$ , as

$$F_i(t, x) = \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} b_i^{(j)}(n, k),$$

and we determine the relations that must be satisfied by the coefficients  $b_i^{(j)}(n, k)$ . We have

$$\begin{aligned} \frac{\partial F_i(t, x)}{\partial t} &= -\lambda F_i(t, x) + \frac{\lambda}{r_j - r_{j-1}^+} \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} \\ &\quad \times [r_j b_i^{(j)}(n + 1, k) - r_{j-1}^+ b_i^{(j)}(n + 1, k + 1)], \end{aligned}$$

and

$$\begin{aligned} \frac{\partial F_i(t, x)}{\partial x} &= \frac{\lambda}{r_j - r_{j-1}^+} \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1 - x_j)^{n-k} \\ &\quad \times [b_i^{(j)}(n + 1, k + 1) - b_i^{(j)}(n + 1, k)]. \end{aligned}$$

Using the uniformization technique, we have

$$\sum_{r \in S} F_r(t, x) A(r, i) = -\lambda F_i(t, x) + \lambda \sum_{r \in S} F_r(t, x) P(r, i),$$

that is,

$$\begin{aligned} & \sum_{r \in S} F_r(t, x) A(r, i) \\ &= -\lambda F_i(t, x) + \lambda \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} \sum_{k=0}^n \binom{n}{k} x_j^k (1-x_j)^{n-k} \sum_{r \in S} b_r^{(j)}(n, k) P(r, i). \end{aligned}$$

It follows that if the  $b_i^{(j)}(n, k)$  are such that

$$(d_i - r_{j-1}^+) b_i^{(j)}(n+1, k+1) + (r_j - d_i) b_i^{(j)}(n+1, k) = (r_j - r_{j-1}^+) \sum_{r \in S} b_r^{(j)}(n, k) P(r, i) \quad (\text{A.1})$$

then Eq. (1) is satisfied.

The recurrence relation (A.1) can also be written as follows, for  $j = u, \dots, m$ .

For  $i \in B_0 \cup \dots \cup B_{j-1}$ ,

$$b_i^{(j)}(n, k) = \frac{r_{j-1}^+ - d_i}{r_j - d_i} b_i^{(j)}(n, k+1) + \frac{r_j - r_{j-1}^+}{r_j - d_i} \sum_{r \in S} b_r^{(j)}(n-1, k) P(r, i)$$

and for  $i \in B_j \cup \dots \cup B_m$ ,

$$b_i^{(j)}(n, k) = \frac{d_i - r_j}{d_i - r_{j-1}^+} b_i^{(j)}(n, k-1) + \frac{r_j - r_{j-1}^+}{d_i - r_{j-1}^+} \sum_{r \in S} b_r^{(j)}(n-1, k-1) P(r, i).$$

Using matrix and vector notation, we get for  $j = u, \dots, m$

$$b_{B_l}^{(j)}(n, k) = \frac{r_l - r_j}{r_l - r_{j-1}^+} b_{B_l}^{(j)}(n, k-1) + \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \sum_{i=0}^m b_{B_i}^{(j)}(n-1, k-1) P_{B_i B_l}$$

for  $0 \leq l \leq j-1$  and

$$b_{B_l}^{(j)}(n, k) = \frac{r_{j-1}^+ - r_l}{r_j - r_l} b_{B_l}^{(j)}(n, k+1) + \frac{r_j - r_{j-1}^+}{r_j - r_l} \sum_{i=0}^m b_{B_i}^{(j)}(n-1, k) P_{B_i B_l}$$

for  $j \leq l \leq m$ .

To get the initial conditions for the  $b_i^{(u)}(n, k)$ , we consider the jumps of  $F_i(t, x)$ .

For  $t > 0$  and  $i \in B_u \cup \dots \cup B_m$ , we have

$$F_i(t, 0) = \Pr\{X_t = i\} = \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^n}{n!} (\alpha P^n)(i),$$

where  $(\alpha P^n)(i)$  denotes the  $i$ th entry of the row vector  $\alpha P^n$ . It follows that

$$b_i^{(u)}(n, 0) = (\alpha P^n)(i),$$

that is

$$b_{B_l}^{(u)}(n, 0) = (\alpha P^n)_{B_l} \quad \text{for } u \leq l \leq m.$$

For  $t > 0$  and  $u \leq j \leq m - 1$  and  $i \notin B_j$ , we have

$$F_i(t, r_j t) = \lim_{x \xrightarrow{\leq} r_j t} F_i(t, x),$$

since  $i \notin B_j$  means that there is no jump at point  $x = r_j t$ . It follows that

$$b_i^{(j+1)}(n, 0) = b_i^{(j)}(n, n) \quad \text{if } i \notin B_j,$$

that is

$$b_{B_l}^{(j+1)}(n, 0) = b_{B_l}^{(j)}(n, n) \quad \text{for } l \neq j.$$

This can also be written as

$$b_{B_l}^{(j)}(n, 0) = b_{B_l}^{(j-1)}(n, n) \quad \text{for } u < j \leq l \leq m,$$

$$b_{B_l}^{(j)}(n, n) = b_{B_l}^{(j+1)}(n, 0) \quad \text{for } 0 \leq l \leq j - 1 < m - 1.$$

Finally, for  $t > 0$  and  $i \notin B_m$ , we have

$$0 = F_i(t, r_m t) = \lim_{x \xrightarrow{\leq} r_m t} F_i(t, x).$$

It follows that  $b_i^{(m)}(n, n) = 0$ , that is

$$b_{B_l}^{(m)}(n, n) = 0 \quad \text{for } 0 \leq l \leq m - 1.$$

The proof is now complete.

### Appendix B. Proof of Theorem 2

The proof is made by successive inductions using the relations described in Theorem 1.

*Step 0.* For  $n = 0$  and for every  $j = u, \dots, m$ , we have

$$b_{B_l}^{(j)}(0, 0) = 0_{B_l} \quad \text{for } 0 \leq l \leq j - 1, \quad b_{B_l}^{(j)}(0, 0) = \alpha_{B_l} \quad \text{for } j \leq l \leq m.$$

So relation (3) is satisfied for  $n = 0$ .

*Step 1.* Suppose relation (3) is satisfied for integer  $n - 1$  and let us prove that it is true for integer  $n \geq 1$ .

Let  $u \leq j \leq m$ .

*Step 1.1.* We first consider the case where  $0 \leq l \leq j - 1$ .

- For  $j = m$  we have from Theorem 1,
  - $b_{B_l}^{(m)}(n, n) = 0_{B_l} \leq (\alpha P^n)_{B_l}$ .
  - Suppose that  $b_{B_l}^{(m)}(n, k + 1) \leq (\alpha P^n)_{B_l}$  for integer  $k \leq n - 1$ . Then

$$b_{B_l}^{(m)}(n, k) = \frac{r_{m-1}^+ - r_l}{r_m - r_l} b_{B_l}^{(m)}(n, k + 1) + \frac{r_m - r_{m-1}^+}{r_m - r_l} \sum_{i=0}^m b_{B_i}^{(m)}(n - 1, k) P_{B_i B_l}$$

$$\begin{aligned} &\leq \frac{r_{m-1}^+ - r_l}{r_m - r_l} (\alpha P^n)_{B_l} + \frac{r_m - r_{m-1}^+}{r_m - r_l} \sum_{i=0}^m (\alpha P^{n-1})_{B_i} P_{B_i B_l} \\ &= \frac{r_{m-1}^+ - r_l}{r_m - r_l} (\alpha P^n)_{B_l} + \frac{r_m - r_{m-1}^+}{r_m - r_l} (\alpha P^n)_{B_l} \\ &= (\alpha P^n)_{B_l}. \end{aligned}$$

So the relation is satisfied for  $n \geq 0$ , for  $j = m$ , for  $0 \leq k \leq n$ , and for  $0 \leq l \leq m - 1$ .

- Suppose now that the relation is satisfied for integer  $j + 1$ ,  $j \leq m - 1$ .

Using Theorem 1, we have:

- $b_{B_l}^{(j)}(n, n) = b_{B_l}^{(j+1)}(n, 0) \leq (\alpha P^n)_{B_l}$ .
- Suppose that  $b_{B_l}^{(j)}(n, k + 1) \leq (\alpha P^n)_{B_l}$  for integer  $k \leq n - 1$ . Then

$$\begin{aligned} b_{B_l}^{(j)}(n, k) &= \frac{r_{j-1}^+ - r_l}{r_j - r_l} b_{B_l}^{(j)}(n, k + 1) + \frac{r_j - r_{j-1}^+}{r_j - r_l} \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k) P_{B_i B_l} \\ &\leq \frac{r_{j-1}^+ - r_l}{r_j - r_l} (\alpha P^n)_{B_l} + \frac{r_j - r_{j-1}^+}{r_j - r_l} \sum_{i=0}^m (\alpha P^{n-1})_{B_i} P_{B_i B_l} \\ &= \frac{r_{j-1}^+ - r_l}{r_j - r_l} (\alpha P^n)_{B_l} + \frac{r_j - r_{j-1}^+}{r_j - r_l} (\alpha P^n)_{B_l} \\ &= (\alpha P^n)_{B_l}. \end{aligned}$$

So the relation is satisfied for  $n \geq 0$ , for  $j = u, \dots, m$ , for  $0 \leq k \leq n$ , and for  $0 \leq l \leq j - 1$ .

Step 1.2. In the same way, we now consider the case where  $j \leq l \leq m$ .

- For  $j = u$  we have from Theorem 1:

- $b_{B_l}^{(u)}(n, 0) = (\alpha P^n)_{B_l}$ .
- Suppose that  $b_{B_l}^{(m)}(n, k - 1) \leq (\alpha P^n)_{B_l}$  for integer  $k \geq 1$ . Then

$$\begin{aligned} b_{B_l}^{(u)}(n, k) &= \frac{r_l - r_u}{r_l} b_{B_l}^{(u)}(n, k - 1) + \frac{r_u}{r_l} \sum_{i=0}^m b_{B_i}^{(u)}(n - 1, k - 1) P_{B_i B_l} \\ &\leq \frac{r_l - r_u}{r_l} (\alpha P^n)_{B_l} + \frac{r_u}{r_l} \sum_{i=0}^m (\alpha P^{n-1})_{B_i} P_{B_i B_l} \\ &= \frac{r_l - r_u}{r_l} (\alpha P^n)_{B_l} + \frac{r_u}{r_l} (\alpha P^n)_{B_l} \\ &= (\alpha P^n)_{B_l}. \end{aligned}$$

So the relation is satisfied for  $n \geq 0$ , for  $j = u$ , for  $0 \leq k \leq n$ , and for  $u \leq l \leq m$ .

- Suppose now that the relation is satisfied for integer  $j - 1$ ,  $j \geq u + 1$ .

Using Theorem 1, we have:

- $b_{B_l}^{(j)}(n, 0) = b_{B_l}^{(j-1)}(n, n) \leq (\alpha P^n)_{B_l}$ .

- Suppose that  $b_{B_l}^{(j)}(n, k - 1) \leq (\alpha P^n)_{B_l}$  for integer  $k \geq 1$ . Then

$$\begin{aligned} b_{B_l}^{(j)}(n, k) &= \frac{r_l - r_j}{r_l - r_{j-1}} b_{B_l}^{(j)}(n, k - 1) + \frac{r_j - r_{j-1}}{r_l - r_{j-1}} \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k - 1) P_{B_i B_l} \\ &\leq \frac{r_l - r_j}{r_l - r_{j-1}} (\alpha P^n)_{B_l} + \frac{r_j - r_{j-1}}{r_l - r_{j-1}} \sum_{i=0}^m (\alpha P^{n-1})_{B_i} P_{B_i B_l} \\ &= \frac{r_l - r_j}{r_l - r_{j-1}} (\alpha P^n)_{B_l} + \frac{r_j - r_{j-1}}{r_l - r_{j-1}} (\alpha P^n)_{B_l} \\ &= (\alpha P^n)_{B_l}. \end{aligned}$$

So the relation is satisfied for  $n \geq 0$ , for  $j = u, \dots, m$ , for  $0 \leq k \leq n$ , and for  $j \leq l \leq m$ , which completes the proof.

### Appendix C. Proof of Theorem 3

The proof of relation (4) is immediate since, for  $j > u$ , we have

$$b_{B_l}^{(j)}(n, 0) = b_{B_l}^{(j-1)}(n, n) - \alpha_{B_{j-1}} P_{B_{j-1} B_{j-1}}^n \mathbf{1}_{B_{j-1}} \mathbf{1}_{\{l=j-1\}},$$

where  $\mathbf{1}_{\{c\}} = 1$  if condition  $c$  is satisfied and 0 otherwise. The proof of relation (5) is made by successive inductions using the relations described in Theorem 1. Note that from Theorem 1,  $b_{B_l}^{(j)}(n, k)$  is a convex combination of two terms. It follows that

$$b_{B_l}^{(j)}(n, k) \leq b_{B_l}^{(j)}(n, k - 1) \Leftrightarrow \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k - 1) P_{B_i B_l} \leq b_{B_l}^{(j)}(n, k) \leq b_{B_l}^{(j)}(n, k - 1)$$

for  $j \leq l \leq m$  and

$$b_{B_l}^{(j)}(n, k) \geq b_{B_l}^{(j)}(n, k + 1) \Leftrightarrow b_{B_l}^{(j)}(n, k + 1) \leq b_{B_l}^{(j)}(n, k) \leq \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, k) P_{B_i B_l}$$

for  $0 \leq l \leq j - 1$ .

*Step 0.* We prove the relation for  $n = 1$ . For  $n = 1$ ,  $j = u$  and  $u \leq l \leq m$  we have from Theorem 2

$$b_{B_l}^{(u)}(1, 1) \leq (\alpha P)_{B_l} = b_{B_l}^{(u)}(1, 0).$$

Suppose that the relation is satisfied at level  $j - 1$ ,  $j < m$ . Then for  $j \leq l \leq m$ , we have from Theorem 1 and using the equivalence above

$$\begin{aligned} b_{B_l}^{(j)}(1, 0) - b_{B_l}^{(j)}(1, 1) &= \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \left[ b_{B_l}^{(j)}(1, 0) - \sum_{i=0}^m b_{B_i}^{(j)}(0, 0) P_{B_i B_l} \right] \\ &\geq \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \left[ b_{B_l}^{(j-1)}(1, 1) - \sum_{i=0}^m b_{B_i}^{(j-1)}(0, 0) P_{B_i B_l} \right] \geq 0. \end{aligned}$$

So the relation is satisfied for  $n = 1$  and  $j \leq l \leq m$ .



For  $n = 1$ ,  $j = m$  and  $0 \leq l \leq m - 1$  we have

$$b_{B_l}^{(m)}(1, 1) = 0 \leq b_{B_l}^{(m)}(1, 0).$$

Suppose that the relation is satisfied at level  $j + 1$ ,  $j > u$ . Then for  $0 \leq l \leq j - 1$ , we have from Theorem 1 and using the equivalence above

$$\begin{aligned} b_{B_l}^{(j)}(1, 0) - b_{B_l}^{(j)}(1, 1) &= \frac{r_j - r_{j-1}^+}{r_j - r_l} \left[ \sum_{i=0}^m b_{B_i}^{(j)}(0, 0) P_{B_i B_l} - b_{B_l}^{(j)}(1, 1) \right] \\ &\geq \frac{r_j - r_{j-1}^+}{r_j - r_l} \left[ \sum_{i=0}^m b_{B_i}^{(j+1)}(0, 0) P_{B_i B_l} - b_{B_l}^{(j+1)}(1, 0) \right] \geq 0. \end{aligned}$$

So the relation is satisfied for  $n = 1$  and  $0 \leq l \leq j - 1$ .

Step 1. Suppose relation (5) is satisfied for integer  $n - 1$  and let us prove that it is true for integer  $n$ ,  $n \geq 2$ . Let  $u \leq j \leq m$ .

Step 1.1. We first consider the case where  $0 \leq l \leq j - 1$ .

- For  $j = m$  we have from Theorem 1:
  - $b_{B_l}^{(m)}(n, n) = 0_{B_l} \leq b_{B_l}^{(m)}(n, n - 1)$ .
  - Suppose that  $b_{B_l}^{(m)}(n, k + 1) \geq b_{B_l}^{(m)}(n, k + 2)$  for integer  $k \leq n - 2$ . Then

$$\begin{aligned} &b_{B_l}^{(m)}(n, k) - b_{B_l}^{(m)}(n, k + 1) \\ &= \frac{r_{m-1}^+ - r_l}{r_m - r_l} [b_{B_l}^{(m)}(n, k + 1) - b_{B_l}^{(m)}(n, k + 2)] \\ &\quad + \frac{r_m - r_{m-1}^+}{r_m - r_l} \sum_{i=0}^m [b_{B_i}^{(m)}(n - 1, k) - b_{B_i}^{(m)}(n - 1, k + 1)] P_{B_i B_l}, \end{aligned}$$

which shows that  $b_{B_l}^{(m)}(n, k) - b_{B_l}^{(m)}(n, k + 1) \geq 0$ .

So the relation is satisfied for  $n \geq 1$ , for  $j = m$ , for  $1 \leq k \leq n$ , and for  $0 \leq l \leq m - 1$ .

- Suppose now that the relation is satisfied at level  $j + 1$ ,  $j \leq m - 1$ .
  - Using Theorem 1, we have

$$\begin{aligned} b_{B_l}^{(j)}(n, n - 1) - b_{B_l}^{(j)}(n, n) &= \frac{r_j - r_{j-1}^+}{r_j - r_l} \left[ \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, n - 1) - b_{B_l}^{(j)}(n, n) \right] \\ &\geq \frac{r_j - r_{j-1}^+}{r_j - r_l} \left[ \sum_{i=0}^m b_{B_i}^{(j+1)}(n - 1, 0) - b_{B_l}^{(j+1)}(n, 0) \right] \\ &\geq 0. \end{aligned}$$

- Suppose that  $b_{B_l}^{(j)}(n, k + 1) \geq b_{B_l}^{(j)}(n, k + 2)$  for integer  $k \leq n - 2$ . Then,

$$b_{B_l}^{(j)}(n, k) - b_{B_l}^{(j)}(n, k + 1)$$

$$\begin{aligned}
 &= \frac{r_{j-1}^+ - r_l}{r_j - r_l} [b_{B_l}^{(j)}(n, k + 1) - b_{B_l}^{(j)}(n, k + 2)] \\
 &\quad + \frac{r_j - r_{j-1}^+}{r_j - r_l} \sum_{i=0}^m [b_{B_i}^{(j)}(n - 1, k) - b_{B_i}^{(j)}(n - 1, k + 1)] P_{B_i B_l},
 \end{aligned}$$

which shows that  $b_{B_l}^{(j)}(n, k) - b_{B_l}^{(j)}(n, k + 1) \geq 0$ .

So the relation is satisfied for  $n \geq 1$ , for  $j = u, \dots, m$ , for  $1 \leq k \leq n$ , and for  $0 \leq l \leq j - 1$ .

Step 1.2. In the same way, we now consider the case where  $j \leq l \leq m$ .

• For  $j = u$  we have from Theorem 3:

- $b_{B_l}^{(u)}(n, 0) = (\alpha P^n)_{B_l} \geq b_{B_l}^{(u)}(n, 1)$ .
- Suppose that  $b_{B_l}^{(u)}(n, k - 2) - b_{B_l}^{(u)}(n, k - 1) \geq 0$ , for integer  $k \geq 2$ . Then

$$\begin{aligned}
 &b_{B_l}^{(u)}(n, k - 1) - b_{B_l}^{(u)}(n, k) \\
 &= \frac{r_l - r_u}{r_l} [b_{B_l}^{(u)}(n, k - 2) - b_{B_l}^{(u)}(n, k - 1)] \\
 &\quad + \frac{r_u}{r_l} \sum_{i=0}^m [b_{B_i}^{(u)}(n - 1, k - 2) - b_{B_i}^{(u)}(n - 1, k - 1)] P_{B_i B_l},
 \end{aligned}$$

which shows that  $b_{B_l}^{(u)}(n, k - 1) - b_{B_l}^{(u)}(n, k) \geq 0$ .

So the relation is satisfied for  $n \geq 1$ , for  $j = u$ , for  $1 \leq k \leq n$ , and for  $u \leq l \leq m$ .

• Suppose now that the relation is satisfied for integer  $j - 1$ ,  $j \geq u + 1$ .

• Using Theorem 1, we have

$$\begin{aligned}
 &b_{B_l}^{(j)}(n, 0) - b_{B_l}^{(j)}(n, 1) \\
 &= \frac{r_j - r_{j-1}}{r_l - r_{j-1}} \left[ b_{B_l}^{(j)}(n, 0) - \sum_{i=0}^m b_{B_i}^{(j)}(n - 1, 0) P_{B_i B_l} \right] \\
 &\geq \frac{r_j - r_{j-1}}{r_l - r_{j-1}} \left[ b_{B_l}^{(j-1)}(n, n) - \sum_{i=0}^m b_{B_i}^{(j-1)}(n - 1, n - 1) P_{B_i B_l} \right] \geq 0.
 \end{aligned}$$

• Suppose that  $b_{B_l}^{(j)}(n, k - 2) - b_{B_l}^{(j)}(n, k - 1) \geq 0$ , for integer  $k \geq 2$ . Then,

$$\begin{aligned}
 &b_{B_l}^{(j)}(n, k - 1) - b_{B_l}^{(j)}(n, k) \\
 &= \frac{r_l - r_j}{r_l - r_{j-1}^+} [b_{B_l}^{(j)}(n, k - 2) - b_{B_l}^{(j)}(n, k - 1)] \\
 &\quad + \frac{r_j - r_{j-1}^+}{r_l - r_{j-1}^+} \sum_{i=0}^m [b_{B_i}^{(j)}(n - 1, k - 2) - b_{B_i}^{(j)}(n - 1, k - 1)] P_{B_i B_l},
 \end{aligned}$$

which shows that  $b_{B_l}^{(j)}(n, k - 1) - b_{B_l}^{(j)}(n, k) \geq 0$ .

So the relation is satisfied for  $n \geq 1$ , for  $j = u, \dots, m$ , for  $1 \leq k \leq n$ , and for  $j \leq l \leq m$ , which completes the proof.

## References

- [1] D. Anick, D. Mitra, M.M. Sondhi, Stochastic theory of a data-handling system with multiple sources, *Bell System Tech. J.* 61 (8) (1982) 1871–1894.
- [2] A. Dupuis, F. Guillemin, B. Sericola, Asymptotic results for the superposition of a large number of data connections on an ATM link, *Queueing Systems* 26 (1–2) (1997).
- [3] F. Guillemin, G. Rubino, B. Sericola, A. Simonian, Transient analysis of statistical multiplexing of data connections on an ATM link, in: *ITC'15 – 15th International Teletraffic Congress*, Washington, DC, USA, June 1997.
- [4] B. Igel'nik, Y. Kogan, V. Krivan, D. Mitra, A new computational approach for stochastic fluid models of multiplexers with heterogeneous sources, *Queueing Systems* 20 (1995) 85–116.
- [5] H. Kobayashi, Q. Ren, A mathematical theory for transient analysis of communication networks, *IEICE Trans. Comm.* E75-B(12) (1992) 1266–1276.
- [6] L. Kosten, Stochastic theory of data-handling systems with groups of multiple sources, in: *Proc. IFIP WG 7.3/TC 6 Second International Symposium on the Performance of Computer-Communication Systems*, Zurich, Switzerland, March 1984, pp. 321–331.
- [7] D. Mitra, Stochastic fluid models, in: *Proc. Performance'87*, Brussels, Belgium, December 1987, pp. 39–51.
- [8] D. Mitra, Stochastic theory of a fluid model of producers and consumers coupled by a buffer, *Adv. Appl. Probab.* 20 (1988) 646–676.
- [9] H. Nabli, B. Sericola, Performability analysis: A new algorithm, *IEEE Trans. Comput.* 45 (4) (1996) 491–494.
- [10] A. Narayanan, V.G. Kulkarni, First passages times in fluid models with an application to two priority fluid systems, in: *Proc. IPDS'96*, Urbana – Champaign, IL, USA, September 1996, pp. 166–175.
- [11] Q. Ren, H. Kobayashi, Transient solutions for the buffer behavior in statistical multiplexing, *Performance Evaluation* 23 (1995) 65–87.
- [12] J.W. Roberts (Ed.), *Performance Evaluation and Design of Multiservice Networks*, Commission of the European Communities, Information Technologies and Sciences, 1992, Final report of the COST 224 Project.
- [13] J.W. Roberts, U. Mocci, J. Virtamo (Eds.), *Broadband Network Teletraffic. Performance Evaluation and Design of Broadband Multiservice Networks*, Lecture Notes in Computer Science, vol. 1155, Springer, Berlin, 1995, Final report of the COST 242 action.
- [14] T.E. Stern, A.I. Elwalid, Analysis of separable Markov-modulated rate models for information-handling systems, *Adv. Appl. Probab.* 23 (1991) 105–139.
- [15] T. Tanaka, O. Hashida, Y. Takahashi, Transient analysis of fluid models for ATM statistical multiplexer, *Performance Evaluation* 23 (1995) 145–162.



**Bruno Sericola** received the Ph.D. degree in computer science from the University of Rennes I in 1988. He has been with INRIA (Institut National de Recherche en Informatique et Automatique) since 1989. His main research activity is in performance and dependability analysis of computer and communication systems, in particular using Markov models.